

Is the η – η' complex an ordinary two-level system?★

D. Klabučar^a (speaker), D. Horvatić^a

★ ACHT 2015 – Austria-Croatia-Hungary Triangle workshop
Leibnitz, Austria, 7.– 9. October 2015.

^aPhysics Department, University of Zagreb, Croatia

Introduction

QUESTION: do η and η' always obey von Neumann–Wigner **anticrossing theorem**, stating that *if a Hermitian matrix represents an observable for a system and depends on continuous real parameters, its eigenvalues cannot cross (i.e., its eigenvalues cannot become exactly equal) as any of the parameters vary.*

In the case of a two-state (two-level) system, it is easy to see the avoided crossing of the levels. Namely, the eigenvalues of the 2x2 Hamiltonian matrix \hat{H} are

$$E_{\pm} = \frac{1}{2}(H_{11} + H_{22}) \pm \frac{1}{2}\sqrt{(H_{11} - H_{22})^2 + 4|H_{12}|^2}.$$

Thus E_{\pm} obey $E_{+} > E_{-}$ for all parameter values if the transition matrix element $H_{12} \neq 0$.

\Rightarrow If they form a two-level system, any description of η and η' through their 2x2 mass matrix should also exhibit this property regarding their masses.

However, functional renormalization group approach in a quark-meson truncation indicates that the assignment $M_{\eta'} > M_{\eta}$ changes as the $U_A(1)$ breaking is turned off. The mass assignments in the η - η' complex are thus re-examined.

The η - η' complex in the pseudoscalar nonet

- Pseudoscalar mesons of light quarks $q = u, d, s$ are (almost) Goldstone bosons of DChSB, so for $m_{u,d,s} \rightarrow 0$ also vanishing meson masses² $M_{u\bar{d}}^2 = M_{\pi^+}^2, M_{u\bar{s}}^2 = M_K^2, \dots, \hat{M}_{NA}^2 = \text{diag}(M_{u\bar{u}}^2, M_{d\bar{d}}^2, M_{s\bar{s}}^2)$
QCD chiral behavior reproduced correctly by Dyson-Schwinger-Bethe-Salpeter approach (DS) – except anomalously heavy η' !
- $|u\bar{d}\rangle = |\pi^+\rangle, |u\bar{s}\rangle = |K^+\rangle, \dots$ but $|u\bar{u}\rangle, |d\bar{d}\rangle$ and $|s\bar{s}\rangle$ do not correspond to any physical particles (at $T = 0$ at least!), although in the isospin limit (adopted from now on) $M_{u\bar{u}} = M_{d\bar{d}} = M_{u\bar{d}} = M_{\pi}$. $I = \text{good}$ Q.no. \Rightarrow recouple into "more physical" $I_3 = 0$ octet-singlet basis


$$I = 1 \quad |\pi^0\rangle = \frac{1}{\sqrt{2}}(|u\bar{u}\rangle - |d\bar{d}\rangle),$$

$$\text{but } I = 0 \quad |\eta_8\rangle = \frac{1}{\sqrt{6}}(|u\bar{u}\rangle + |d\bar{d}\rangle - 2|s\bar{s}\rangle) \approx |\eta\rangle \quad \text{mixes with}$$

$$I = 0 \quad |\eta_0\rangle = \frac{1}{\sqrt{3}}(|u\bar{u}\rangle + |d\bar{d}\rangle + |s\bar{s}\rangle) \approx |\eta'\rangle \quad \dots \text{too heavy for GB}.$$

Except η - η' , pseudoscalars qualitatively understood at both $T = 0$ and $T > 0$

- An illustration of 'non-anomalous' meson $M_s(T)$ by a simple 'separable' DS model:



- 'Deconfinement' $T_{d,q}$ from S_q pole - very different $T_{d,u}$, $T_{d,s}$... can be cured/synchronized with $T_{Ch}(= T_{cri})$ by **Polyakov loop**
- But what about η and η' both at $T = 0$ and $T > 0$?

Physical η and η' must have a diagonal mass matrix

- the “non-anomalous” (**chiral-limit-vanishing!**) part of the mass-squared matrix of π^0 and η 's is (in π^0 - η_8 - η_0 basis)

$$\hat{M}_{NA}^2 = \begin{pmatrix} M_\pi^2 & 0 & 0 \\ 0 & M_{88}^2 & M_{80}^2 \\ 0 & M_{08}^2 & M_{00}^2 \end{pmatrix} \xrightarrow[\text{U}_A(1) \text{ problem}]{\text{diagonalization}} \begin{pmatrix} M_\pi^2 & 0 & 0 \\ 0 & M_\pi^2 & 0 \\ 0 & 0 & M_{s\bar{s}}^2 \end{pmatrix}$$

$$M_{88}^2 \equiv \langle \eta_8 | \hat{M}_{NA}^2 | \eta_8 \rangle = \frac{2}{3} (M_{s\bar{s}}^2 + \frac{1}{2} M_\pi^2), \quad M_{00}^2 \equiv \langle \eta_0 | \hat{M}_{NA}^2 | \eta_0 \rangle = \frac{2}{3} (\frac{1}{2} M_{s\bar{s}}^2 + M_\pi^2),$$

$$M_{80}^2 \equiv \langle \eta_8 | \hat{M}_{NA}^2 | \eta_0 \rangle = M_{08}^2 = \frac{\sqrt{2}}{3} (M_\pi^2 - M_{s\bar{s}}^2)$$

- What reproduces M_π & M_K cannot also $M_\eta = 548$ & $M_{\eta'} = 958$ MeV!
- \hat{M}_{NA}^2 **not enough!** To avoid **the $U_A(1)$ problem**, one must break the $U_A(1)$ symmetry (**as it is destroyed by the gluon anomaly**) at least at the level of the masses.

Why $\eta_0 \approx \eta'$ has an anomalous piece of mass:

$U_A(1)$ symmetry is broken by nonabelian ("gluon") axial anomaly: **even in the chiral limit** (ChLim, where $m_q \rightarrow 0$),

$$\partial_\alpha \bar{\psi}(x) \gamma^\alpha \gamma_5 \frac{\lambda^0}{2} \psi(x) \propto F^a(x) \cdot \tilde{F}^a(x) \equiv \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu}^a(x) F_{\rho\sigma}^a(x) \neq 0.$$

This breaks the $U_A(1)$ symmetry of QCD and precludes the 9^{th} Goldstone pseudoscalar meson \Rightarrow very massive η' : **even in ChLim**, where $m_\pi, m_K, m_\eta \rightarrow 0$, **still ('ChLim WVR')**

$$0 \neq \Delta M_{\eta_0}^2 = \Delta M_{\eta'}^2 = \frac{(A = \text{qty. dim. mass})^4}{(“f_{\eta'}”)^2} = \frac{6 \chi_{\text{YM}}}{f_\pi^2} + O\left(\frac{1}{N_c}\right)$$

Out of ChLim : $M_{\eta'}^2 + M_\eta^2 - 2 M_K^2 = \frac{2N_f}{f_\pi^2} \chi_{\text{YM}} \left(+O\left(\frac{1}{N_c}\right) \right)$

Anomalous part of η_0 mass: $\Delta M_{\eta_0}^2 = \chi_{\text{YM}} \frac{2N_f}{f_\pi^2} + O\left(\frac{1}{N_c}\right)$

QCD chiral behavior (reproduced by DS approach) **of the non-anomalous parts** of masses of light $q\bar{q}'$ pseudoscalars (i.e., all parts except ΔM_{η_0}): $M_{q\bar{q}'}^2 = \text{const} (m_q + m_{q'})$, ($q, q' = u, d, s$).

\Rightarrow non-anomalous parts of the masses in WVR cancel:

$$M_{\eta'}^2 + M_\eta^2 - 2M_K^2 \approx \Delta M_{\eta_0}^2, \quad \text{approx. as in ChLim WVR}$$

$$\chi = \int d^4x \langle 0|Q(x)Q(0)|0\rangle, \quad Q(x) = \frac{g^2}{64\pi^2} \epsilon_{\mu\nu\rho\sigma} F_{\mu\nu}^a(x) F_{\rho\sigma}^a(x)$$

- $Q(x)$ = topological charge density operator
- In WV rel., χ is the pure-gluon, YM one, $\chi_{\text{YM}} \leftrightarrow \chi_{\text{quench}}$.

Lattice: good χ_{YM} , subtleties with χ of light-flavor QCD [Bernard et al.,

JHEP 1206 (2012) 051] where
$$\chi = - \frac{\langle \bar{q}q \rangle_0}{\sum_{q=u,d,s} \frac{1}{m_q}} + \mathcal{C}(\text{higher } \mathcal{O} \text{ in } m).$$

Gluon anomaly is not accessible to ladder approximation


- All masses in $\hat{M}_{N_A}^2$ are calculated in the ladder approx., which cannot include the gluon anomaly contributions.
- Large N_c : the gluon anomaly suppressed as $1/N_c!$ → Include its effect just at the level of masses: break the $U_A(1)$ symmetry and avoid the $U_A(1)$ problem by shifting the η_0 (squared) mass by anomalous contribution 3β .
- complete mass matrix is then $\hat{M}^2 = \hat{M}_{N_A}^2 + \hat{M}_A^2$ where

$$\hat{M}_A^2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 3\beta \end{pmatrix} \quad \text{does not vanish in the chiral limit.}$$

$3\beta = \Delta M_{\eta_0}^2$ = the anomalous mass² of η_0 [in SU(3) limit incl. ChLim] is **related to the YM topological susceptibility**. Fixed by phenomenology or (here) **taken from the lattice**.

Transitions related to the $U_A(1)$ anomaly

- Transitions between hidden flavors $|q\bar{q}\rangle \rightarrow |q'\bar{q}'\rangle$
($q, q' = u, d, s$)



- Diamond graph: **just the simplest example** of a transition $|q\bar{q}\rangle \rightarrow |q'\bar{q}'\rangle$
($q, q' = u, d, s$), contributing to the anomalous masses in the η - η' complex, but not included in the interaction kernel in the ladder approximation.

Anomalous mass matrix in $q\bar{q}$ and octet-singlet bases

- we can also rewrite \hat{M}_A^2 in the $q\bar{q}$ basis $|u\bar{u}\rangle, |d\bar{d}\rangle, |s\bar{s}\rangle$

$$\hat{M}_A^2 = \beta \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \xrightarrow[\text{breaking}]{\text{flavor}} \hat{M}_A^2 = \beta \begin{pmatrix} 1 & 1 & X \\ 1 & 1 & X \\ X & X & X^2 \end{pmatrix}$$

- We introduced the **effects of the flavor breaking** on the anomaly-induced transitions $|q\bar{q}\rangle \rightarrow |q'\bar{q}'\rangle$ ($q, q' = u, d, s$). $s\bar{s}$ transition suppression estimated by $X \approx f_\pi / f_{s\bar{s}}$.
- Then, \hat{M}_A^2 in the octet-singlet basis is modified to

$$\hat{M}_A^2 = \beta \begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{2}{3}(1-X)^2 & \frac{\sqrt{2}}{3}(2-X-X^2) \\ 0 & \frac{\sqrt{2}}{3}(2-X-X^2) & \frac{1}{3}(2+X)^2 \end{pmatrix}$$

- \rightarrow **In the isospin limit**, one can always restrict to 2×2 submatrix of etas ($I=0$), as π^0 ($I=1$) **is decoupled then**.

Anomalous mass matrix and mixing in NS – S basis

- nonstrange (NS) – strange (S) basis

$$|\eta_{NS}\rangle = \frac{1}{\sqrt{2}}(|u\bar{u}\rangle + |d\bar{d}\rangle) = \frac{1}{\sqrt{3}}|\eta_8\rangle + \sqrt{\frac{2}{3}}|\eta_0\rangle,$$

$$|\eta_S\rangle = |s\bar{s}\rangle = -\sqrt{\frac{2}{3}}|\eta_8\rangle + \frac{1}{\sqrt{3}}|\eta_0\rangle.$$

- the η – η' mass matrix in this basis is

$$\hat{M}^2 = \begin{pmatrix} M_{\eta_{NS}}^2 & M_{\eta_S\eta_{NS}}^2 \\ M_{\eta_{NS}\eta_S}^2 & M_{\eta_S}^2 \end{pmatrix} = \begin{pmatrix} M_{u\bar{u}}^2 + 2\beta & \sqrt{2}\beta X \\ \sqrt{2}\beta X & M_{s\bar{s}}^2 + \beta X^2 \end{pmatrix} \xrightarrow{\phi} \begin{pmatrix} M_{\eta}^2 & 0 \\ 0 & M_{\eta'}^2 \end{pmatrix}$$

- NS – S mixing relations – states rotation diagonalizing \hat{M}^2 :

$$|\eta\rangle = \cos\phi|\eta_{NS}\rangle - \sin\phi|\eta_S\rangle, \quad |\eta'\rangle = \sin\phi|\eta_{NS}\rangle + \cos\phi|\eta_S\rangle.$$

$$\theta = \phi - \arctan\sqrt{2}$$

Finally, fix anomalous contribution to η - η' :

- Equal traces of diagonalized & non-diagonalized. \hat{M}^2 demand 1st equality in

$$\beta(2+X^2) = M_\eta^2 + M_{\eta'}^2 - 2M_K^2 = \frac{2N_f}{f_\pi^2} \chi_{\text{YM}} \quad (2^{\text{nd}} \text{ equality} = \text{WV rel.})$$

- requiring that the experimental trace $(M_\eta^2 + M_{\eta'}^2)_{\text{exp}} \approx 1.22 \text{ GeV}^2$ be reproduced by the theoretical \hat{M}^2 , yields

$$\beta_{\text{fit}} = \frac{1}{2+X^2} [(M_\eta^2 + M_{\eta'}^2)_{\text{exp}} - (M_{u\bar{u}}^2 + M_{s\bar{s}}^2)]$$
- Or, get β from lattice χ_{YM} ! Then no free parameters!
- then, can calculate the NS - S mixing angle ϕ

$$\tan 2\phi = \frac{2 M_{\eta_S \eta_{NS}}^2}{M_{\eta_S}^2 - M_{\eta_{NS}}^2} = \frac{2 \sqrt{2} \beta X}{M_{\eta_S}^2 - M_{\eta_{NS}}^2} \quad \text{and}$$

$$M_{\eta_{NS}}^2 = M_{u\bar{u}}^2 + 2\beta = M_\pi^2 + 2\beta, \quad M_{\eta_S}^2 = M_{s\bar{s}}^2 + \beta X^2 = M_{s\bar{s}}^2 + \beta \frac{f_\pi^2}{f_{s\bar{s}}^2}$$

Physical η, η' eigenmasses – of the two-level type:

- The diagonalization of the NS - S mass matrix then finally gives us the *calculated* η and η' masses:

$$M_\eta^2 = \cos^2 \phi M_{\eta_{NS}}^2 - M_{\eta_S \eta_{NS}}^2 \sin 2\phi + \sin^2 \phi M_{\eta_S}^2 \quad (\text{note } M_{\eta_S \eta_{NS}}^2 = \sqrt{2}\beta X)$$

$$M_{\eta'}^2 = \sin^2 \phi M_{\eta_{NS}}^2 + M_{\eta_S \eta_{NS}}^2 \sin 2\phi + \cos^2 \phi M_{\eta_S}^2$$

- Equivalently, secular determinant \Rightarrow the eigenvalues of 2×2 matrix:

$$M_\eta^2 = \frac{1}{2} \left[M_{\eta_{NS}}^2 + M_{\eta_S}^2 - \sqrt{(M_{\eta_{NS}}^2 - M_{\eta_S}^2)^2 + 4 M_{\eta_S \eta_{NS}}^4} \right]$$

$$= \frac{1}{2} \left[M_\pi^2 + M_{s\bar{s}}^2 + \beta(2 + X^2) - \sqrt{(M_\pi^2 + 2\beta - M_{s\bar{s}}^2 - \beta X^2)^2 + 8\beta^2 X^2} \right]$$

$$M_{\eta'}^2 = \frac{1}{2} \left[M_{\eta_{NS}}^2 + M_{\eta_S}^2 + \sqrt{(M_{\eta_{NS}}^2 - M_{\eta_S}^2)^2 + 4 M_{\eta_S \eta_{NS}}^4} \right]$$

$$= \frac{1}{2} \left[M_\pi^2 + M_{s\bar{s}}^2 + \beta(2 + X^2) + \sqrt{(M_\pi^2 + 2\beta - M_{s\bar{s}}^2 - \beta X^2)^2 + 8\beta^2 X^2} \right]$$

Separable model results on η and η' at $T = 0$


	β_{fit}	$\beta_{\text{latt.}}$	Exp.
θ	-12.22°	-13.92°	
M_η [MeV]	548.9	543.1	547.75
$M_{\eta'}$ [MeV]	958.5	932.5	957.78
X	0.772	0.772	
3β [GeV ²]	0.845	0.781	

- $X = f_\pi / f_{s\bar{s}}$ as well as the whole \hat{M}_{NA}^2 (consisting of M_π and $M_{s\bar{s}}$) are calculated model quantities.
- $\beta_{\text{latt.}}$ was obtained from $\chi_{\text{YM}}(T = 0) = (175.7 \text{ MeV})^4$
- **But is an extension to high T possible**, as there is a large mismatch of characteristic temperature scales of the pure-gauge YM ($T_c \sim 270$ MeV) vs. full QCD ($T_c \sim 160$ MeV) with quarks?
- Concretely in WVR, χ_{YM} is more T -resistant than QCD quantities $M_{\eta,\eta',K}$ and f_π . Does WVR become unusable as T approaches the (pseudo-)critical temperatures of full QCD, such as $T \sim T_{\text{Ch}}$?

Assume: $\chi_{\text{YM}}(T)/f_{\pi}^2(T)$ gives the T -dependence of anomalous mass





just to illustrate anticrossing! Experiment nowadays EXCLUDES such modeling of $\Delta M_{\eta_0}(T)$.

$M_{\rho}[\text{GeV}], T_c=128.\text{MeV}$




Assume: $\chi_{\text{YM}}(T)/f_\pi^2(T)$ gives the T -dependence of anomalous mass

just to illustrate anticrossing! Experiment nowadays EXCLUDES such modeling of $\Delta M_{\eta_0}(T)$.



Comparison of pseudoscalars including η_0 and η_8

$M_p[\text{GeV}], T_c=128.\text{MeV}$



Discrepancy with phenomenology removed by another relation connecting YM and QCD!

Shore's generalization of WV valid to all orders in $1/N_c$

- WV rel. – lowest order in $1/N_c$ – improved like this:

$$(f_{\eta'}^0)^2 M_{\eta'}^2 + (f_{\eta}^0)^2 M_{\eta}^2 = \frac{1}{3} (f_{\pi}^2 M_{\pi}^2 + 2f_K^2 M_K^2) + 6A \quad (1)$$

$$f_{\eta'}^0 f_{\eta'}^8 M_{\eta'}^2 + f_{\eta}^0 f_{\eta}^8 M_{\eta}^2 = \frac{2\sqrt{2}}{3} (f_{\pi}^2 M_{\pi}^2 - f_K^2 M_K^2) \quad (2)$$

$$(f_{\eta'}^8)^2 M_{\eta'}^2 + (f_{\eta}^8)^2 M_{\eta}^2 = -\frac{1}{3} (f_{\pi}^2 M_{\pi}^2 - 4f_K^2 M_K^2) \quad (3)$$

A is the full QCD topological charge parameter (replacing χ_{YM} in WV)

$$A = \frac{\chi}{1 + \chi \left(\frac{1}{\langle \bar{u}u \rangle m_u} + \frac{1}{\langle \bar{d}d \rangle m_d} + \frac{1}{\langle \bar{s}s \rangle m_s} \right)} \quad (4)$$

= seemed hard to calculate on lattice (maybe easier today?) ...

However, it is known that $A = \chi_{\text{YM}} + \mathcal{O}\left(\frac{1}{N_c}\right)$ (at $T = 0$)

Approximating the full QCD topological charge parameter A

Replacing **3 different condensates** by the **chiral one**, $\langle \bar{q}q \rangle_0$, reduces the **full QCD topological charge A** (4) to the combination $\tilde{\chi}$ on the RHS of Leutwyler-Smilga relation:

$$\chi_{\text{YM}} = \frac{\chi}{1 + \frac{\chi}{\langle \bar{q}q \rangle_0} \sum_{q=u,d,s} \frac{1}{m_q}} \rightarrow \tilde{\chi}(T, \mu) = \frac{\langle \bar{q}q(T, \mu) \rangle_0}{\sum_{q=u,d,s} \frac{1}{m_q}} + \text{corr}'s \approx A(T, \mu)$$

because of Di Vecchia-Veneziano result for small m_q :

$$\chi = - \frac{m \langle \bar{q}q \rangle_0}{N_f} + \text{corrections}(m),$$

(Previously, we only **conjectured** $\chi_{\text{YM}}(T) \rightarrow \tilde{\chi}(T)$ [Benić& al, Phys.Rev.D84 (2011)016006].)

\Rightarrow The quark condensates $\langle \bar{q}q(T, \mu) \rangle$, and **not the pure-gauge** quantity χ_{YM} , determine the T (and μ) dependence of (partial) restoration of $U_A(1)$. \Rightarrow **Linked with the chiral restoration!**

T -dependence of χ and $\tilde{\chi}$


- Extending the light-quark full-QCD topol. susceptibility χ is somewhat uncertain, as there is no guidance from lattice [unlike for $\chi_{\text{YM}}(T)$].
- The leading term in Di Vecchia-Veneziano relation $\propto \langle \bar{q}q \rangle_0(T)$ very plausibly, but for the correction term we have to explore a range of Ansätze, i.e.,

$$\chi(T) = -\frac{m \langle \bar{q}q \rangle_0(T)}{N_f} + \mathcal{C}(m) \left[\frac{\langle \bar{q}q \rangle_0(T)}{\langle \bar{q}q \rangle_0(T=0)} \right]^\delta, \quad (0 \leq \delta < 2).$$

Then, $\tilde{\chi}(T) =$

$$= \frac{\langle \bar{q}q \rangle_0(T)}{\sum_{q=u,d,s} \left(\frac{1}{m_q} \right)} \left\{ 1 - \frac{\langle \bar{q}q \rangle_0(T)}{\sum_{q=u,d,s} \left(\frac{1}{m_q} \right)} \frac{1}{\mathcal{C}(m)} \left[\frac{\langle \bar{q}q \rangle_0(T=0)}{\langle \bar{q}q \rangle_0(T)} \right]^\delta \right\}.$$


Chiral condensate $\langle q\bar{q} \rangle_0(T)$ and resulting $\tilde{\chi}(T)$



Case 1: T -independent correction term in χ


[Benić, Horvatić, Kekez and Klabučar, Phys. Rev. D 84 (2011) 016006.]

$$\chi_{\text{YM}} = (0.1757 \text{ GeV})^4, \delta = 0$$




Case 2: Strongly T -dependent correction term $\propto \langle \bar{q}q \rangle_0(T)$

$$\chi_{\text{YM}} = (0.1757 \text{ GeV})^4, \delta = 1$$



T -dependence of the NS-S mixing angle $\phi(T)$




$\phi(T)$ for the cases of the T -independent correction term in $\chi(T)$ ($\delta = 0$) and the correction term in $\chi(T)$ behaving like the leading term, i.e., like the chiral condensate ($\delta = 1$), and for two values of $\tilde{\chi}(T = 0) = \chi_{YM}$.

A functional renormalization group (FRG) approach


[M. Mitter & B. J. Schaefer, Phys. Rev. D **89** (2014) 5, 054027]: Axial anomaly & chiral symmetry investigated by a FRG approach in a three flavor quark-meson truncation.

- Chiral order parameters: quark condensates $\langle \bar{q}q \rangle$ related via bosonization to vacuum expectation values of the scalar-isoscalar mesonic fields $\sigma_{x,y} = \sigma_{NS,S}$:

$$\langle \Sigma \rangle = \text{diag}(\langle \sigma_x \rangle / 2, \langle \sigma_x \rangle / 2, \langle \sigma_y \rangle / \sqrt{2})$$



NS (x) & S (y) condensates in the nonstrange chiral limit





With explicit chiral breaking for all flavors

In the FRG approach to the quark-meson truncation, condensates dominate $U_A(1)$ breaking

The $U_A(1)$ breaking implemented in the effective Lagrangian through the 't Hooft determinantal interaction ξ with the coupling strength c :

$$c \xi = c \left(\det [\Sigma] + \det [\Sigma^\dagger] \right) , \text{ where } \Sigma = T^a (\sigma_a + i\pi_a)$$

First panel: $c \neq 0$ contributes to good agreement with the present data:





Second panel: no $U_A(1)$ breaking, due to $c = 0$. η changed little, but η' is now **degenerate with π , i.e., lighter** than η .

Thus, **varying** of the coupling parameter c between 0 and its phenomenological value **would lead to level crossing** for any T .

Linear Σ -model+quarks, Polyakov L.& (axi-)vector mesons

P. Kovacs, Z. Szep and G. Wolf, J. Phys. Conf. Ser. 599 (2015) 1, 012010



Summary

- The mass assignments in the η - η' complex has been re-examined during melting of the axial anomaly. The assignment $M_{\eta'} > M_{\eta}$ holds through (anti)crossings of matrix elements of the η - η' mass matrix.
- The general condition enabling crossings of eigenvalues is that the terms should belong to different irreducible representations of the symmetry group of the Hamiltonian of the system. There is no indication that this may be the case in the η - η' complex. Thus a Hermitean mass matrix contains a complete description of the masses of η and η' mesons. The η - η' complex remains an ordinary two-level system.
- The η - η' mass crossing seemingly exhibited by a functional renormalization group approach in a quark-meson truncation is probably an artifact of this level of truncation (which, however, can be systematically improved).

Additional slides

On Shore's generalization of WV relation
and
its combining with the
Feldmann–Kroll–Stech (FKS) scheme

η' and η have 4 independent decay constants

$$f_{\eta'}^0, f_{\eta}^8, f_{\eta}^0, f_{\eta'}^8 : \quad \langle 0 | A^{a\mu}(x) | P(p) \rangle = i f_P^a p^\mu e^{-ip \cdot x}, \quad a = 8, 0; \quad P = \eta, \eta'.$$

- Equivalently, one has 4 related but different constants $f_{\eta'}^{NS}, f_{\eta}^{NS}, f_{\eta}^S, f_{\eta'}^S$ if instead of octet and singlet axial currents ($a = 8, 0$) one takes this matrix element of the nonstrange-strange axial currents ($a = NS, S$)

$$A_{NS}^\mu(x) = \frac{1}{\sqrt{3}} A^{8\mu}(x) + \sqrt{\frac{2}{3}} A^{0\mu}(x) = \frac{1}{2} (\bar{u}(x) \gamma^\mu \gamma_5 u(x) + \bar{d}(x) \gamma^\mu \gamma_5 d(x)),$$

$$A_S^\mu(x) = -\sqrt{\frac{2}{3}} A^{8\mu}(x) + \frac{1}{\sqrt{3}} A^{0\mu}(x) = \frac{1}{\sqrt{2}} \bar{s}(x) \gamma^\mu \gamma_5 s(x),$$

$$\begin{bmatrix} f_{\eta}^{NS} & f_{\eta}^S \\ f_{\eta'}^{NS} & f_{\eta'}^S \end{bmatrix} = \begin{bmatrix} f_{\eta}^8 & f_{\eta}^0 \\ f_{\eta'}^8 & f_{\eta'}^0 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{3}} & -\sqrt{\frac{2}{3}} \\ \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} \end{bmatrix},$$

$$a, P = NS, S : \quad \langle 0 | A_{NS}^\mu(x) | \eta_{NS}(p) \rangle = i f_{NS} p^\mu e^{-ip \cdot x}, \quad \langle 0 | A_{NS}^\mu(x) | \eta_S(p) \rangle = 0,$$

$$a, P = NS, S : \quad \langle 0 | A_S^\mu(x) | \eta_S(p) \rangle = i f_S p^\mu e^{-ip \cdot x}, \quad \langle 0 | A_S^\mu(x) | \eta_{NS}(p) \rangle = 0,$$

- Note: in a DS approach, $f_{NS} = f_{u\bar{u}} = f_{d\bar{d}} = f_\pi$, $f_S = f_{s\bar{s}}$ are calculated quantities

Two Mixing Angles and FKS one-angle scheme

- Any 4 η - η' decay constants conveniently parametrized in terms of two decay constants and two angles:

$$\begin{aligned}
 f_{\eta}^8 &= \cos \theta_8 f_8, & f_{\eta}^0 &= -\sin \theta_0 f_0, & f_{\eta}^{NS} &= \cos \phi_{NS} f_{NS}, & f_{\eta}^S &= -\sin \phi_S f_S, \\
 f_{\eta'}^8 &= \sin \theta_8 f_8, & f_{\eta'}^0 &= \cos \theta_0 f_0, & f_{\eta'}^{NS} &= \sin \phi_{NS} f_{NS}, & f_{\eta'}^S &= \cos \phi_S f_S
 \end{aligned}$$

- Big **practical** difference between 0-8 and NS - S schemes:
- while θ_8 and θ_0 differ a lot from each other and from $\theta \approx (\theta_8 + \theta_0)/2$, FKS showed that $\phi_{NS} \approx \phi_S \approx \phi$.

$$\begin{bmatrix} f_{\eta}^{NS} & f_{\eta}^S \\ f_{\eta'}^{NS} & f_{\eta'}^S \end{bmatrix} = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} f_{NS} & 0 \\ 0 & f_S \end{bmatrix}.$$

For four decay constants, can use FKS one-angle scheme!

- ϕ relates $\{f_\eta^8, f_{\eta'}^8, f_\eta^0, f_{\eta'}^0\}$ with $\{f_{NS}, f_S\} = \{f_\pi, f_{s\bar{s}}\}$:

$$\begin{bmatrix} f_\eta^8 & f_\eta^0 \\ f_{\eta'}^8 & f_{\eta'}^0 \end{bmatrix} = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} f_{NS} & 0 \\ 0 & f_S \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{3}} & \sqrt{\frac{2}{3}} \\ -\sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} \end{bmatrix}$$

- Some other useful relations between quantities of NS - S (FKS) and 0 - 8 schemes:

$$f_8 = \sqrt{\frac{1}{3}f_{NS}^2 + \frac{2}{3}f_S^2}, \quad \theta_8 = \phi - \arctan\left(\frac{\sqrt{2}f_S}{f_{NS}}\right),$$

$$f_0 = \sqrt{\frac{2}{3}f_{NS}^2 + \frac{1}{3}f_S^2}, \quad \theta_0 = \phi - \arctan\left(\frac{\sqrt{2}f_{NS}}{f_S}\right).$$

Solve numerically Shore's Eqs. (1)-(3) for $M_{\eta'}$, M_{η} , and ϕ :

Inputs:	$M_{\pi}, M_K, f_{\pi} = f_{NS}, f_{s\bar{s}} = f_S$ and f_K , calculated in 3 different DS models					
χ_{YM}	191^4	175.7^4	191^4	175.7^4	191^4	175.7^4
M_{η}	499.8	485.7	496.7	482.8	526.2	507.0
$M_{\eta'}$	931.4	815.8	934.9	818.4	983.2	868.7
ϕ	52.01°	46.11°	51.85°	46.07°	47.23°	40.86°
θ	-2.72°	-8.62°	-2.89°	-8.67°	-7.51°	-13.87°
θ_0	7.74°	1.84°	7.17°	1.39°	-0.33°	-6.69°
θ_8	-12.00°	-17.90°	-11.85°	-17.6°	-14.12°	-20.47°
f_0	108.8	108.8	107.9	107.9	101.8	101.8
f_8	122.6	122.6	121.1	121.1	110.7	110.7
f_{η}^0	-14.7	-3.5	-13.5	-2.6	0.6	11.9
$f_{\eta'}^0$	107.9	108.8	107.1	107.9	101.8	101.1
f_{η}^8	119.9	116.7	118.5	115.4	107.4	103.7
$f_{\eta'}^8$	-25.5	-37.7	-2.49	-37.6	-27.0	-38.7

(in D. Horvatić et al., Eur. Phys. J. A **38** (2008) 257.) $M_{\eta, \eta'}$ and f 's in MeV, χ_{YM} is in MeV^4 .

The same is now reproduced **analytically**:

- Eqs. (1)-(3) \Rightarrow two **closed-form solutions** for M_η , $M_{\eta'}$ and $\tan \phi$ in terms of f_π , $f_{s\bar{s}}$, M_π , M_K and A .

The set reproducing the previous numerical results is:

$$\tan \phi = \frac{-2A f_\pi^2 + 4A f_{s\bar{s}}^2 - 2f_K^2 f_\pi^2 M_K^2 + f_\pi^4 M_\pi^2 + f_\pi^2 f_{s\bar{s}}^2 M_\pi^2 + \Delta}{4\sqrt{2} A f_\pi f_{s\bar{s}}}$$

$$M_{\eta,\eta'}^2 = \frac{2A f_\pi^2 + 4A f_{s\bar{s}}^2 + 2f_K^2 f_\pi^2 M_K^2 - f_\pi^4 M_\pi^2 + f_\pi^2 f_{s\bar{s}}^2 M_\pi^2 \mp \Delta}{2f_\pi^2 f_{s\bar{s}}^2}$$

where $\Delta^2 =$

$$32 A^2 f_\pi^2 f_{s\bar{s}}^2 + \left\{ 2A(f_\pi^2 - 2f_{s\bar{s}}^2) + f_\pi^2 \left[2f_K^2 M_K^2 - (f_\pi^2 + f_{s\bar{s}}^2) M_\pi^2 \right] \right\}^2$$

[Benić, Horvatić, Kekez & Klabučar, Phys. Lett. B738 (2014) 113]

Find matrix elem's in NS - S basis from these $M_\eta, M_{\eta'}, \phi$:

$$\begin{aligned} M_{\eta_{NS}}^2 &\equiv M_{NS}^2 &= \cos^2 \phi M_\eta^2 + \sin^2 \phi M_{\eta'}^2 \\ M_{\eta_S}^2 &\equiv M_S^2 &= \sin^2 \phi M_\eta^2 + \cos^2 \phi M_{\eta'}^2 \\ M_{\eta_{NS}\eta_S}^2 &\equiv M_{NSS}^2 &= \sin \phi \cos \phi (M_\eta^2 - M_{\eta'}^2) \end{aligned}$$

to use
$$M_{\eta, \eta'}^2 = \frac{1}{2} \left[M_{NS}^2 + M_S^2 \mp \sqrt{(M_{NS}^2 - M_S^2)^2 + 4M_{NSS}^2} \right]$$

Mathematica leads to surprisingly simple results:

$$M_{NS}^2 = M_\pi^2 + \frac{4A}{f_\pi^2}, \quad M_{NSS}^2 = \frac{2\sqrt{2}A}{f_\pi f_{s\bar{s}}}$$

$$M_S^2 = \frac{1}{f_{s\bar{s}}^2} [2 f_K^2 M_K^2 - f_\pi^2 M_\pi^2] + \frac{2A}{f_{s\bar{s}}^2} = M_{s\bar{s}}^2 + \frac{2A}{f_{s\bar{s}}^2}$$

$$\begin{aligned} f_\pi^2 M_\pi^2 &= -m_u \langle u\bar{u} \rangle - m_d \langle d\bar{d} \rangle \quad \text{and} \quad f_K^2 M_K^2 = -m_u \langle u\bar{u} \rangle - m_s \langle s\bar{s} \rangle \\ \Rightarrow \quad 2 f_K^2 M_K^2 - f_\pi^2 M_\pi^2 &= f_{s\bar{s}}^2 M_{s\bar{s}}^2 \quad \text{"eq. (23)"} \end{aligned}$$

Compare M_{NS} , M_{NSS} and M_{S} with NS-S mass matrix:

$$\begin{bmatrix} M_{\text{NS}}^2 & M_{\text{NSS}}^2 \\ M_{\text{NSS}}^2 & M_{\text{S}}^2 \end{bmatrix} = \begin{bmatrix} M_{\pi}^2 + 2\beta & \sqrt{2}\beta X \\ \sqrt{2}\beta X & M_{s\bar{s}}^2 + \beta X^2 \end{bmatrix}$$

⇒ Very similar formulas in WV case and "Shore case":

$$1.) \quad \beta_{\text{WV}} = \frac{6\chi_{YM}}{f_{\pi}^2(2 + X^2)}, \quad \beta_{\text{Shore+FKS}} = \frac{2A}{f_{\pi}^2} \approx \frac{2\chi_{YM}}{f_{\pi}^2}$$

Explains why Shore's scheme needs higher values of χ_{YM} than WV, to approach empirical masses.

$$2.) \quad X = \frac{f_{\pi}}{f_{s\bar{s}}} \quad \text{the SAME in the both WV and Shore cases ...}$$

... but in the "Shore case", it follows from equations! Before, incl. WV, it was an input – estimate, educated guess.

T -dependence of pseudoscalar decay constants

